

Original Paper

# Acceptability of a Digital Care App in Patients Undergoing Hip and Knee Arthroplasty: Prospective Cohort Study

Yacine Louni<sup>1</sup>, MDCM; Matthew Laroche<sup>1</sup>, BCom; Abdulrhman Alnasser<sup>2</sup>, MD; Mohammad Abuhaneya<sup>2</sup>, MD; Eric Belzile<sup>3</sup>, MSc; Sandhya Baskaran<sup>3</sup>, MSc; Jennifer Mutch<sup>2</sup>, MDCM, FRCSC; Anthony Albers<sup>2</sup>, MDCM, FRCSC

<sup>1</sup>Faculty of Medicine, McGill University, Montreal, QC, Canada

<sup>2</sup>Division of Orthopaedic Surgery, McGill University, Montreal, QC, Canada

<sup>3</sup>St Mary's Hospital Research Center, St Mary's Hospital, Montreal, QC, Canada

**Corresponding Author:**

Anthony Albers, MDCM, FRCSC  
Division of Orthopaedic Surgery  
McGill University  
St Mary's Hospital, 3830 Avenue Lacombe  
Montreal, QC H3T 0A2  
Canada  
Phone: +1 514-345-3511 ext 3282  
Email: [anthony.albers@mail.mcgill.ca](mailto:anthony.albers@mail.mcgill.ca)

## Abstract

**Background:** Mobile health (mHealth) apps have become more commonly used in orthopedics. However, for these apps to be efficient, patients should be willing to use them, making it essential to understand patients' perspectives of mHealth interventions.

**Objective:** The aim of this single-center, intent-to-treat, preoperative single-cohort study of 100 patients was to evaluate the acceptability of mymobility (Zimmer-Biomet), an mHealth app designed for the postoperative care of total hip arthroplasty (THA) and total knee arthroplasty (TKA).

**Methods:** We measured acceptability using the theoretical framework for acceptability (TFA) preoperatively and at 3 months post operation. We also measured satisfaction with app use postoperatively using the Usefulness, Satisfaction, and Ease of Use questionnaire as well as patient-reported outcome measures preoperatively and postoperatively using the Oxford hip and knee scores and the visual analog scale for pain. Patients included were 18 years or older; underwent unilateral primary total hip, total knee, or partial knee arthroplasty; spoke and read French or English; and had a smartphone with internet access. Participants used mymobility in addition to standard government-funded physiotherapy.

**Results:** The preoperative overall TFA result was 4.2 out of 5, but results decreased significantly postoperatively. There was higher self-efficacy in preoperative TFAs with higher education, and lower acceptability in postoperative TFAs with TKA. The Usefulness, Satisfaction, and Ease of Use questionnaire revealed a good level of satisfaction with the use of the app. Patient-reported outcome measures showed earlier improvement in THA (31.2 d) than in TKA (89.4 d), whereas the visual analog scale showed a rapid decrease in pain with both procedures. Only 1 patient expressed privacy concerns with the use of the app.

**Conclusions:** There was a good level of acceptability with the use of mymobility for the postoperative management of THA and TKA, although acceptability decreased postoperatively. This decrease could signify high expectations toward the app preoperatively or higher than expected difficulty and pain in the early postoperative period. Acceptability tended to increase with higher education and decrease with TKA. These trends are consistent with prior literature and constitute a potential gap to address for app developers. The influence of the natural recovery process on acceptability remains unclear. Future studies could explore this gap by comparing results in cohorts using the app to cohorts with standard care.

**JMIR Hum Factors** 2026;13:e79682; doi: [10.2196/79682](https://doi.org/10.2196/79682)

**Keywords:** mobile health; orthopedics; patient satisfaction; telemedicine; telerehabilitation; total joint arthroplasty

## Introduction

In Canada, there were 60 705 hip and 70 379 knee replacements performed in the inpatient setting in 2023-2024 [1]. These procedures were the third and second most common inpatient surgeries, and these numbers are expected to at least double in the United States (and likely Canada) over the next 15 years due to an aging population [2,3]. Another growing sector in health care is the use of smartphones for patient care. An estimated 71% of the world population uses a smartphone in 2024, and there are more than 350 000 mobile health (mHealth) apps currently available [4,5]. This increasing use of mHealth apps is also present in various areas of orthopedics, where numerous studies have shown that they are efficient and convenient, that they promote patient adherence and engagement with their treatment, and that they facilitate rehabilitation while promoting positive outcomes [6-13].

Although mHealth has the potential to improve outcomes, patients' attitudes toward these apps must be assessed, as they can only be effective if patients are willing to use them. User acceptability is defined as "a multifaceted construct that reflects the extent to which people delivering or receiving health care intervention consider it to be appropriate based on anticipated or experienced cognitive and emotional responses to the intervention" [14]. Acceptability is an important measure, as it not only provides information on patients' experience with mHealth apps, but also gives an insight into their cognitive and emotional response [14]. This helps determine whether patients genuinely perceive them as appropriate and valuable.

Recent studies investigated patients' perspectives of mHealth interventions in the context of postoperative hip and knee arthroplasty care. These studies found a high level of engagement and increased compliance with treatment exercises when using mHealth interventions [13,15-18]. Other findings include participants being satisfied with these interventions while also finding them easy to use and engaging, and providing better connection to their treating team [13,16,19,20]. These results are promising and could signify that patients are willing to use mHealth apps for their rehabilitation. However, these studies either did not assess acceptability or did so through surveys that were researcher-created or not primarily designed to measure acceptability. According to Perski et al [21], this reduces clarity regarding which aspects of acceptability are satisfied, and which are not. Additionally, high engagement does not necessarily reflect patients' perceptions of the intervention. Engagement might be driven by external factors, such as study participation, financial incentives, or free access to the devices and apps, while patients might still find the app to be time-consuming, repetitive, and burdensome.

Measuring acceptability, especially preoperatively, is therefore relevant as it provides a better perspective of patients' opinions prior to and beyond use, and because it "may usefully be considered an emergent property [...] which in turn influences (and is influenced by) user engagement and intervention effectiveness" [21].

We started using the mymobility app (Zimmer Biomet, Warsaw, and Indiana) in 2022 in our community hospital to provide postoperative care to patients receiving outpatient total hip arthroplasty (THA) and total knee arthroplasty (TKA) [22]. This app, available on iOS and Android, currently stands at more than 50 000 downloads on the Google Play Store and is amongst the most downloaded when compared to other apps with the same objective [23]. In this context, our team sought to investigate the acceptability of mymobility, an mHealth app specifically designed for the postoperative care of THA and TKA, using a standardized framework.

## Methods

### Study Design

The primary aim of this single-center, intent-to-treat (ITT), single-cohort study was to evaluate pre- and postoperative patient acceptability with the use of the mymobility app for the postoperative care of THA and TKA. The secondary outcomes were postoperative patient satisfaction with the app and pre- and postoperative patient-reported outcome measures (PROMs).

### Data Collection

Data collected for this study are presented in [Multimedia Appendix 1](#). Demographic data were obtained from patient charts. PROMs were collected either directly through the mymobility app or by one of the authors contacting participants by phone or email. Details of the specific PROMs are provided later in this section. All data were stored securely and deidentified prior to analysis.

### Ethical Considerations

This study received ethics approval from the Research Ethics Board of the Montreal West Island's Integrated University Health and Social Services Centers (Biomedical Subcommittee), affiliated with St. Mary's Hospital, Montreal (IRB# 2024-954). All participants provided informed consent prior to participation. Study data, including personal information, remained confidential and were deidentified. No secondary analyses were conducted. Study participants received no financial compensation, but access to the mymobility app was given for free. No identifiable images of participants were included in this manuscript. Under the Personal Information Protection and Electronic Documents ACT, the data collected by mymobility could not be used or shared with third parties, nor could cookies be collected, unless participants gave their explicit consent within the app [24,25]. If consent was given, data could be used or shared by the app following anonymization, with consent revocable at any time. Participants were not required to give consent to data use or sharing, or to cookie collection to use the app and be part of the study. This was made clear to participants as part of their informed consent.

### App Description

Mymobility is an app that is accessible either through a smartphone or an Apple Watch. The app features educational

content on pre- and postoperative care, exercise routines with metrics to track progress, AI-assisted gait speed analysis to determine the level of recovery, in-app PROMs, and telemedicine with the opportunity to send SMS text messages, videos, or pictures to health care providers (see [Multimedia Appendix 2](#)). The app engages patients in their recovery by sending notifications and reminders if users accept them while also keeping track of progress made. It also provides patients with basic and important smartphone functions to use (eg, Wi-Fi) for those with low digital literacy.

## Questionnaires

### Acceptability

Acceptability was measured using the theoretical framework for acceptability (TFA) questionnaire developed by Sekhon et al [26], which was adapted to our research question (see [Multimedia Appendix 3](#)). The questionnaire comprises 8 items answered by participants preoperatively and at 3 months post operation by phone or email. Each statement was evaluated using a 5-point Likert scale: scores 1 to 2 were classified as negative, 3 as neutral, and 4-5 as positive. The mean score was calculated for each of the 8 items, as well as an overall score.

### Satisfaction

Patient satisfaction was measured using the Usefulness, Satisfaction, and Ease of Use (USE) questionnaire at 3 months post operation (see [Multimedia Appendix 4](#)) [27]. Satisfaction was defined as “the net feeling of pleasure or displeasure that results from aggregating all the benefits that a person hopes to receive from interaction with the information system” [27]. The questionnaire was answered by phone or email. Statements were evaluated using a 7-point Likert scale: scores 1 to 3 were classified as negative, 4 as neutral, and 5 to 7 as positive. The mean score was then calculated for each of the 4 sections of the questionnaire. Two additional questions assessing satisfaction were asked and scored on a 5-point Likert scale (see [Multimedia Appendix 5](#)).

### Patient-Reported Outcome Measures

PROMs were measured using the Oxford Hip Score (OHS) and the Oxford Knee Score (OKS) as well as the visual analog scale (VAS) for pain (scored from 0 to 10) preoperatively and at 1, 3, and 6 months post operation (see [Multimedia Appendices 6 and 7](#)) [28-30]. OHS and OKS were answered directly within the app, whereas the VAS was answered by phone or email.

### Recruitment

Patients were included in the study if they were booked for unilateral primary THA or TKA, were clinically suited for

telehealth care (characterized as age  $\geq 18$  y, speak and read French or English, and have a smartphone or a tablet), had internet access, and provided informed consent to participate in the study. Patients were excluded if they did not meet the inclusion criteria, if they were unable or unwilling to provide informed consent, if they were undergoing emergency THA or TKA, or if they were undergoing revision surgeries. Participants had access to the app in addition to normal postoperative care, which included physiotherapy and postoperative follow-up visits with an orthopedic surgeon.

Patients were approached by a member of the research team during their preoperative assessment visit in the clinic. Eligible patients were provided with an informed consent form detailing the study. Once enrolled, a 1-year free access to the mymobility app was provided to the participants by the treating physician or a member of the research team, after which an email with instructions to download the app was sent. Participants were then required to download the app and agree to the terms and conditions to use it [24].

In the end, 100 participants were included in this study and had a profile created in the app. Participants who did not download the app preoperatively were sent a reminder email and given a few days following surgery, after which they were considered as nondownloaders. In the end, 75 participants downloaded the app and 25 did not. Only participants who downloaded the app were provided with the study questionnaires, as the goal was to assess acceptability, satisfaction, and PROMs with use of the app. Patients who did not download the app remained in the study in the context of the ITT format. However, since deciding not to download the app could signify low acceptability, we contacted participants who did not download the app by phone or email to ask the reason behind their decision.


### Data Analysis

Study data was analyzed using the ITT principle. Data analysis was conducted and reported using STATA 17.0 and R software. Statistical significance was defined as a  $P$  value  $<.05$ .

## Results

The recruitment process is summarized in [Figure 1](#). Baseline characteristics of our population can be found in [Table 1](#).

**Figure 1.** Study enrollment, participation, and questionnaire distribution flow diagram. PROM: patient-reported outcome measure; TFA: theoretical framework for acceptability; USE: Usefulness, Satisfaction, and Ease of Use.



**Table 1.** Baseline demographic and clinical characteristics of the study population.

Baseline variables	Participants (N=100)
Female, n (%)	51 (52)
Missing	1 (1)
Age (years), mean (SD)	67.0 (10.7)
Ethnicity, n (%)	
Caucasian	75 (82)
Asian	9 (10)
Arab	3 (3)
Other	5 (5)
Missing	8 (8)
Education, n (%)	
Graduate	20 (21)
Undergraduate	27 (29)
Postsecondary (nonuniversity)	25 (27)
Primary/secondary	22 (23)
Missing	6 (6)

Baseline variables	Participants (N=100)
Employment, n (%)	
Full/part time	31 (34)
Retired	56 (61)
Unemployed	5 (5)
Missing	8 (8)
Language used, n (%)	
English	64 (69)
French	23 (25)
Both	6 (6)
Missing	7 (7)
Comorbidities, n (%)	
Yes <sup>a</sup>	39 (39)
None	61 (61)
Hypertension	18 (31)
Diabetes	10 (17)
Obesity	6 (10)
Thyroid	4 (7)
Cholesterol	4 (7)
Other	17 (29)
Procedure, n (%)	
TKA <sup>b</sup>	55 (55)
THA <sup>c</sup>	42 (42)
PKA <sup>d</sup> /UKA <sup>e</sup>	3 (3)
Income (CAD \$ <sup>f</sup> ), n (%)	
<25,000	8 (9)
25,000-50,000	15 (17)
50,000-75,000	10 (11)
>100,000	24 (27)
Prefer not to say/unknown	13 (15)
Missing	11 (11)
Area of residency, n (%)	
Urban	72 (77)
Rural	22 (23)
Missing	6 (6)

<sup>a</sup>Participants who answered “Yes” can have multiple comorbidities.

<sup>b</sup>TKA: total knee arthroplasty.

<sup>c</sup>THA: total hip arthroplasty.

<sup>d</sup>PKA: partial knee arthroplasty.



<sup>e</sup>UKA: unicompartmental knee arthroplasty.

<sup>f</sup>A currency exchange rate of CAD \$1=US \$0.75 is applicable.

## Acceptability

The overall score for the preoperative TFA questionnaires was 4.2 (SD 0.6; [Table 2](#)). We found a statistically significant increase in self-efficacy with a university level of education compared to non-university ( $P=.04$ ), but no difference related to age, sex, or employment ([Figure 2](#)). There was a statistically significant decrease between pre- and postoperative overall TFA ( $P=.007$ ), as well as in pre- and postoperative

perceived effectiveness ( $P=.01$ ) and self-efficacy ( $P=.008$ ; [Table 3](#)). Furthermore, when looking at TFA results by procedure, there was a statistically significant decrease with TKA in multiple TFA items, including overall acceptability ( $P=.008$ ; [Figure 3](#)). There were no statistically significant differences between pre- and postoperative overall TFA nor in any of the TFA items with THA.

**Figure 2.** Preoperative TFA results (all items) for the overall population and by education level. TFA: theoretical framework for acceptability.**Figure 3.** Comparison between pre- and postoperative overall TFA results for the overall population and by procedure. TFA: theoretical framework for acceptability; THR: total hip replacement; TKR: total knee replacement.

**Table 2.** Preoperative TFA<sup>a</sup> results and subgroup analysis by demographic category<sup>b</sup>.

TFA items	Overall		Age <70 (n=34), mean (SD)	≥70 (n=29), mean (SD)	P value ( <i>t</i> test)	Sex Male (n=33), mean (SD)	Female (n=30), mean (SD)	P value ( <i>t</i> test)	Education Nonuniversity (n=28), mean (SD)	University (n=32), mean (SD)	P value ( <i>t</i> test)	Not working (n=36), mean (SD)	Working (n=23), mean (SD)	P value ( <i>t</i> test)
	(n=63), mean (SD)	(n=63), mean (SD)												
Affective attitude	4.3 (0.9)	4.3 (0.8)	4.4 (0.8)	.84	4.4 (0.8)	4.1 (1.0)	.21	4.1 (1.0)	.21	4.5 (0.8)	.08	4.4 (0.8)	4.3 (1.1)	.67
Burden	3.9 (0.9)	3.9 (1.0)	3.8 (0.9)	.63	4.0 (0.7)	3.9 (1.1)	.47	3.7 (1.2)	.47	4.0 (0.7)	.20	3.8 (1.0)	4.0 (0.9)	.63
Ethicality	4.2 (1.0)	4.1 (1.1)	4.3 (1.1)	.32	4.1 (1.1)	4.3 (0.8)	.34	4.3 (0.9)	.34	4.1 (1.1)	.54	4.1 (1.1)	4.4 (0.8)	.22
Perceived effectiveness	4.2 (0.8)	4.1 (1.0)	4.3 (0.6)	.23	4.2 (0.9)	4.2 (0.9)	.96	4.2 (0.8)	.96	4.2 (0.9)	.98	4.2 (0.7)	4.3 (1.2)	.55
Intervention coherence	4.0 (0.9)	3.8 (1.1)	4.3 (0.6)	.05	4.0 (0.9)	4.0 (1.0)	.99	4.1 (0.7)	.99	3.9 (1.1)	.49	4.1 (0.8)	3.8 (1.2)	.21
Self-efficacy	4.3 (0.7)	4.3 (0.9)	4.3 (0.5)	.93	4.3 (0.8)	4.3 (0.7)	.99	4.1 (0.9)	.99	4.5 (0.6)	.04	4.3 (0.5)	4.3 (1.1)	.89
Opportunity	4.3 (0.9)	4.2 (1.0)	4.5 (0.7)	.16	4.3 (0.9)	4.3 (0.8)	.88	4.4 (0.7)	.88	4.3 (0.9)	.40	4.4 (0.7)	4.2 (1.0)	.27
General acceptability	4.4 (0.5)	4.3 (0.6)	4.4 (0.5)	.74	4.3 (0.5)	4.4 (0.6)	.87	4.3 (0.5)	.87	4.5 (0.5)	.09	4.3 (0.5)	4.5 (0.6)	.40
Overall score <sup>c</sup>	4.2 (0.6)	4.1 (0.6)	4.3 (0.5)	.22	4.2 (0.5)	4.2 (0.7)	.78	4.1 (0.6)	.78	4.2 (0.5)	.55	4.2 (0.5)	4.2 (0.6)	.94

<sup>a</sup>TFA: theoretical framework for acceptability.<sup>b</sup>Statistically significant results are given in italic.<sup>c</sup>Burden, ethicality, and opportunity costs were reversed to compute the overall TFA score.

**Table 3.** Change in TFA<sup>a</sup> from preoperative to 3 months post operation<sup>b</sup>.

TFA items	Overall (n=49)			TKA <sup>c</sup> (n=27)			THA <sup>d</sup> (n=22)		
	Pre, mean (SD)	Post, mean (SD)	P value (t test)	Pre, mean (SD)	Post, mean (SD)	P value (t test)	Pre, mean (SD)	Post, mean (SD)	P value (t test)
Affective attitude	4.5 (0.6)	4.2 (1.3)	.09	4.5 (0.6)	4.1 (1.3)	.22	4.6 (0.7)	4.3 (1.3)	.26
Burden	4.0 (0.8)	4.0 (1.1)	.91	3.9 (1.0)	3.8 (1.3)	.70	4.1 (0.7)	4.3 (0.8)	.36
Ethicality	4.3 (0.9)	3.9 (1.3)	.11	4.4 (0.9)	3.6 (1.4)	.04	4.2 (0.9)	4.3 (1.1)	.89
Perceived effectiveness	4.3 (0.8)	3.8 (1.3)	.01	4.3 (0.7)	3.7 (1.4)	.02	4.2 (0.9)	4.0 (1.2)	.28
Intervention coherence	4.1 (0.9)	3.8 (1.2)	.09	4.2 (0.8)	3.6 (1.3)	.01	3.9 (1.1)	4.0 (1.0)	.69
Self-efficacy	4.4 (0.5)	4.0 (1.2)	.008	4.4 (0.5)	3.8 (1.4)	.02	4.5 (0.6)	4.3 (0.9)	.21
Opportunity	4.4 (0.7)	4.1 (1.1)	.08	4.4 (0.7)	4.0 (1.3)	.10	4.4 (0.8)	4.3 (0.7)	.54
General acceptability	4.4 (0.5)	4.3 (1.0)	.56	4.4 (0.5)	4.1 (1.1)	.28	4.4 (0.5)	4.5 (0.8)	.63
Overall score <sup>e</sup>	4.3 (0.5)	4.0 (0.8)	.007	4.3 (0.5)	3.8 (0.9)	.008	4.3 (0.4)	4.2 (0.6)	.41

<sup>a</sup>TFA: theoretical framework for acceptability.<sup>b</sup>Statistically significant results are given in italic.<sup>c</sup>TKA: total knee arthroplasty.<sup>d</sup>THA: total hip arthroplasty.<sup>e</sup>Burden, ethicality, and opportunity costs were reversed to compute the overall TFA score.

## Satisfaction

The USE questionnaire and the additional satisfaction questions revealed good levels of satisfaction with the use of

the app (Table 4 and Figure 4). Patients consistently rated the app as useful and easy to integrate into their recovery process.


**Table 4.** Usefulness, Satisfaction, and Ease of Use questionnaire results 3 months post operation.

Variable	Participants (n=51)
Usefulness score (1-7)	
Disagree (1-3), n (%)	9 (18)
Neutral (4), n (%)	4 (8)
Agree (5-7), n (%)	37 (74)
Missing, n (%)	1 (1.9)
Mean (SD)	5.0 (1.5)
Satisfaction (1-7)	
Disagree (1-3), n (%)	11 (22)
Neutral (4), n (%)	3 (6)
Agree (5-7), n (%)	36 (72)
Missing, n (%)	1 (1.9)
Mean (SD)	4.9 (1.5)
Ease of use (1-7)	
Disagree (1-3), n (%)	3 (6)
Neutral (4), n (%)	9 (18)
Agree (5-7), n (%)	38 (76)
Missing, n (%)	1 (1.9)
Mean (SD)	5.4 (1.1)
Ease of learning (1-7)	
Disagree (1-3), n (%)	2 (4)
Neutral (4), n (%)	0 (0)
Agree (5-7), n (%)	48 (96)
Missing, n (%)	1 (1.9)
Mean (SD)	5.9 (0.9)

Variable	Participants (n=51)
Additional questions	
Level of satisfaction with postoperative care (1-5)	
Disagree (1-2), n (%)	6 (12)
Neutral (3), n (%)	4 (8)
Agree (4-5), n (%)	41 (80)
Mean (SD)	4.1 (1.2)
Likelihood to recommend care to friend or family member (1-5)	
Disagree (1-2), n (%)	8 (16)
Neutral (3), n (%)	3 (6)
Agree (4-5), n (%)	40 (78)
Mean (SD)	4.2 (1.3)


**Figure 4.** Satisfaction questionnaire results: (A) USE questionnaire results on a stacked bar chart for Likert data; (B) additional satisfaction question results on a 100% stacked bar chart. USE: Usefulness, Satisfaction, and Ease of Use.

#### USE questionnaire results



(A)

#### Additional satisfaction questions



(B)

### Patient-Reported Outcome Measures

There was a statistically significant increase in overall Oxford score and OHS at 31.2 days post operation, but only at 89.4


days post operation in OKS (Table 5 and Figure 5). As for the VAS, there was a significant decrease in overall, hip, and knee VAS scores at 31.2 days post operation.

**Table 5.** Oxford scores and VAS<sup>a</sup> results at 1, 3, and 6 months post operation<sup>b</sup>.

Sample time point	Preop <sup>c</sup>		First postop (with preop data) <sup>d</sup>				Second postop (with preop data) <sup>e</sup>				Third postop (with preop data) <sup>f</sup>			
	Participant s, n	Mea n (SD)	Participant s, n	Mea n (SD)	Mean difference	P value	Participant s, n	Mea n (SD)	Mean difference	P value	Participant s, n	Mea n (SD)	Mean difference	P value
					e vs pre (t test)	e vs pre (t test)	(t test)							
Oxford score overall (0-48)	49	24.1 (10.1)	49	30.4 (8.6)	6.3	<.001	21	37.8 (5.7)	10.4	<.001	9	39.1 (3.8)	12.2	.002
Oxford Hip Score (0-48)	20	22.8 (11.8)	20	35.0 (6.4)	12.3	<.001	8	40.5 (4.4)	13.1	.01	1	— <sup>g</sup>	—	—
Oxford Knee Score (0-48)	29	25.0 (8.9)	29	27.2 (8.6)	2.3	.17	13	36.1 (5.9)	8.7	.001	8	39.9 (3.3)	12.5	.004
VAS overall (0-10)	49	4.6 (2.6)	49	2.3 (2.1)	-2.3	<.001	22	0.9 (1.1)	-3.1	<.001	9	1.1 (0.4)	-2.4	.007
VAS hip (0-10)	20	4.5 (2.6)	20	1.5 (1.4)	-3.0	<.001	9	0.7 (0.3)	-2.7	.009	1	—	—	—
VAS knee (0-10)	29	4.7 (2.6)	29	2.9 (2.2)	-1.8	<.001	13	1.0 (0.4)	-3.4	<.001	8	0.8 (0.3)	-2.7	.004

<sup>a</sup>VAS: visual analog scale.<sup>b</sup>Statistically significant results are given in italic.<sup>c</sup>Gap with surgery (days): mean 18.8 (SD 10.1, range 3-36).<sup>d</sup>Gap with surgery (days): mean 31.2 (SD 2.8, range 30-44).<sup>e</sup>Gap with surgery (days): mean 89.4 (SD 5.1, range 67-93).<sup>f</sup>Gap with surgery (days): mean 180.1 (SD 0.4).<sup>g</sup>Not available.

**Figure 5.** Pre- and postoperative (1, 3, and 6 mo) patient-reported outcome measure results by procedure: (A) Oxford scores and (B) VAS scores. THR: total hip replacement; TKR: total knee replacement; VAS: visual analog scale.



## App Nondownload Rate

Of the 25 participants who did not download the app, 7 provided a reason. Their responses included technological concerns ( $n=2$ ), feeling preoccupied and anxious preparing for the procedure ( $n=1$ ), family issues ( $n=2$ ), missing download instructions ( $n=1$ ), and privacy concerns ( $n=1$ ). The remaining 18 participants were also contacted on multiple occasions but never responded.

## Discussion

### Principal Findings

In this study, we investigated the acceptability of the mymobility app for the postoperative care of 100 participants undergoing THA and TKA. We sought to investigate acceptability using the TFA to ensure all aspects of acceptability were measured.

Preoperatively, overall acceptability was satisfactory, with all but one TFA item scoring above 4 out of 5. The sole exception was burden (3.9/5), which may reflect lower familiarity with smartphone apps among older participants in our sample. We also found that higher education was associated with a higher level of self-efficacy. While general and overall acceptability were not significantly higher with

higher education, this trend aligns with Wang et al's [15], Nuovo et al's [31], and Lee et al's [32] results, who described trends of higher engagement and adherence with higher education and lower receptivity to digital health technologies with lower education. Thus, these findings reveal the importance for app developers to consider this gap and tailor their content to diverse populations, notably through the inclusion of images or videos, which are more easily accessible than heavy-text content.

When comparing pre- and postoperative TFAs, although overall scores were satisfactory at both time points, we found a significant decrease in overall acceptability postoperatively. We suspect this decrease is attributed to elevated expectations toward the app, which were adjusted following its use. However, we believe acceptability could also have been influenced by the natural process of recovery. Indeed, the decline in TFA was significant in TKA, but not in THA. PROM results showed the same trend, with no improvement until 3 months post operation in TKA as opposed to 1 month post operation in THA. This is consistent with Booth et al's [19] and Bourne et al's [33] results showing that TKA patients felt a higher need for in-person rehabilitation as opposed to THA and that short-term outcomes are superior in THA than in TKA. When considering the TFA items that decreased in TKA, which are linked to the perceived helpfulness of the app, this pushes us to believe

that the decrease in postoperative acceptability was likely also driven by the surgical outcomes. This illustrates the possibility for app developers to better address the challenges in TKA rehabilitation, notably through adjustment of patient expectations following this procedure.

When looking at participants who did not download the app, reasons included concerns regarding technology in 2 cases, which might be due to the age of the participants (77-79 y). Additionally, 1 participant expressed privacy concerns. Although we explained how data collection and privacy functioned within the app, we still expected this number to be higher. This result is a positive sign for patient trust toward platforms used to provide care, although the need to discuss patient confidentiality within the app remains important.

## Comparison With Previous Work

Multiple studies have investigated patient perspectives on the use of mHealth apps for joint arthroplasty. These studies all reported a high level of engagement, adherence, and satisfaction [13,15-19,31]. Factors that influenced the results positively included higher education and supportive environments, whereas those that did negatively included higher age, lower economic status, and worsened physical or psychological condition [15,17,31]. Patients also described better health care accessibility and reduced isolation with these interventions [20]. Our results showed that higher education was associated with greater acceptability. However, unlike prior work, we did not observe other demographic characteristics to significantly influence acceptability.

## Strengths and Limitations

The main strength of this study is the measure of acceptability using a validated tool (TFA), which allowed us to assess all domains of this concept. Using this framework, we measured acceptability preoperatively, prior to exposure to the app, and postoperatively, after using the app. This allowed us to interpret how preexposure expectations were met following use of the app. Other strengths include the use of a single app, which reduced heterogeneity; the ITT design, which reflects real-world adherence; and the diverse population.

## Acknowledgments

The authors sincerely thank St. Mary's Hospital for its support. The authors declare that no generative artificial intelligence was used in the preparation and writing of this article. All authors reviewed the results, approved the final version of the manuscript, and agreed to be accountable for all aspects of the work.

## Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## Data Availability

All data generated or analyzed during this study are included in this published article (and the multimedia appendices).

## Conflicts of Interest

A Albers is a paid consultant for Zimmer-Biomet. The other authors declare no conflicts of interest.

## Multimedia Appendix 1

Overview of data collected by the research team and by the mymobility app.

[[DOCX File \(Microsoft Word File, 16 KB\)](#)-Multimedia Appendix 1]

This study also has limitations. The main limitation is the absence of a control group, which limits the ability to attribute the results specifically to the app. This limitation is a result of ethical concerns brought by our institutional review board, which prohibited us from having a control group that could be denied a beneficial intervention. We also observed a high rate of unfilled questionnaires, which could have created a response bias. We believe this is due to most questionnaires being distributed by email or phone, creating a limiting step in communication. Other important limitations include the use of a single app, limiting the generalizability of the results; a potential selection bias toward patients willing to use the app; and statistical analysis being only performed with *t* tests, which might have limited the power of the conclusions. In addition, there was no assessment of baseline technological literacy, which could have affected results despite the app's technological guidance features.

## Conclusion

There was a good level of acceptability with the use of mymobility for the postoperative management of THA and TKA, although acceptability decreased postoperatively. This decrease could signify high expectations toward the app preoperatively or higher than expected difficulty and pain in the early postoperative period. Higher education was associated with higher preoperative acceptability. TKA was associated with lower postoperative acceptability, which could be related to the delayed recovery in TKA when compared to THA. These 2 trends have previously been described in the literature, indicating potential gaps to address for app developers. We found a good level of satisfaction with the app, and Oxford scores showed better outcomes in THA than TKA, whereas VAS showed equally significant decreases in pain with both procedures. Privacy concerns were rare, with only 1 patient reporting them.

The impact of the natural recovery process in total joint arthroplasty on the acceptability of mHealth remains unclear. This gap could be addressed through a comparison of acceptability pre- and postoperatively in cohorts using the app and cohorts with standard care.

## Multimedia Appendix 2

Overview of the mymobility functions and interface. (A) Features of the app with depictions of the app interface and (B) Walk-AI, the mymobility artificial intelligence–assisted gait speed analysis predicting and determining level of recovery based on gait speed.

[[DOCX File \(Microsoft Word File\), 641 KB-Multimedia Appendix 2](#)]

## Multimedia Appendix 3

Modified theoretical framework for acceptability questionnaire.

[[DOC File \(Microsoft Word File\), 52 KB-Multimedia Appendix 3](#)]

## Multimedia Appendix 4

Usefulness, Satisfaction, and Ease of Use questionnaire.

[[DOC File \(Microsoft Word File\), 37 KB-Multimedia Appendix 4](#)]

## Multimedia Appendix 5

Additional satisfaction questions.

[[DOC File \(Microsoft Word File\), 34 KB-Multimedia Appendix 5](#)]

## Multimedia Appendix 6

Oxford Hip and Oxford Knee Scores.

[[DOC File \(Microsoft Word File\), 83 KB-Multimedia Appendix 6](#)]

## Multimedia Appendix 7

Visual analog scale for pain.

[[DOC File \(Microsoft Word File\), 269 KB-Multimedia Appendix 7](#)]

## References

1. Hospital stays in Canada, 2021–2022. Canadian Institute for Health Information; Feb 23, 2023. URL: <https://bit.ly/3N6xCNM> [Accessed 2025-12-09]
2. Shichman I, Roof M, Askew N, et al. Projections and epidemiology of primary hip and knee arthroplasty in Medicare patients to 2040-2060. JB JS Open Access. 2023;8(1):e22.00112. [doi: [10.2106/JBJS.OA.22.00112](https://doi.org/10.2106/JBJS.OA.22.00112)] [Medline: [36864906](https://pubmed.ncbi.nlm.nih.gov/36864906/)]
3. Aging and chronic diseases: a profile of Canadian seniors. Public Health Agency of Canada; 2020. URL: <https://www.canada.ca/en/public-health/services/publications/diseases-conditions/aging-chronic-diseases-profile-canadian-seniors-report.html> [Accessed 2025-12-09]
4. Sherif A. Global smartphone penetration rate as share of population from 2016 to 2024. Statista. 2025. URL: <https://www.statista.com/statistics/203734/global-smartphone-penetration-per-capita-since-2005/> [Accessed 2025-12-09]
5. Wells C, Spry C. An overview of smartphone apps: CADTH horizon scan. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health; 2022. URL: <https://www.ncbi.nlm.nih.gov/books/NBK595384> [Accessed 2025-12-09]
6. Andrawis JP, Muzykewicz DA, Franko OI. Mobile device trends in orthopedic surgery: rapid change and future implications. Orthopedics. 2016;39(1):e51-6. [doi: [10.3928/01477447-20151228-01](https://doi.org/10.3928/01477447-20151228-01)] [Medline: [26730684](https://pubmed.ncbi.nlm.nih.gov/26730684/)]
7. Dionisi S, Giannetta N, Di Simone E, et al. The use of mHealth in orthopedic surgery: a scoping review. Int J Environ Res Public Health. Nov 28, 2021;18(23):12549. [doi: [10.3390/ijerph182312549](https://doi.org/10.3390/ijerph182312549)] [Medline: [34886274](https://pubmed.ncbi.nlm.nih.gov/34886274/)]
8. Hayat J, Ramadhan M, Gonnah AR, Alfadhl A, Al-Naseem AO. The role of mobile health technology in perioperative spinal care: a systematic scoping review and narrative synthesis. Cureus. Feb 2024;16(2):e54254. [doi: [10.7759/cureus.54254](https://doi.org/10.7759/cureus.54254)] [Medline: [38496189](https://pubmed.ncbi.nlm.nih.gov/38496189/)]
9. Mapinduzi J, Ndacyisaba G, Verbrugghe J, Timmermans A, Kossi O, Bonnechère B. Effectiveness of mHealth interventions to improve pain intensity and functional disability in individuals with hip or knee osteoarthritis: a systematic review and meta-analysis. Arch Phys Med Rehabil. Feb 2025;106(2):280-291. [doi: [10.1016/j.apmr.2024.06.008](https://doi.org/10.1016/j.apmr.2024.06.008)] [Medline: [38945508](https://pubmed.ncbi.nlm.nih.gov/38945508/)]
10. Pritwani S, Shrivastava P, Pandey S, et al. Mobile and computer-based applications for rehabilitation monitoring and self-management after knee arthroplasty: scoping review. JMIR Mhealth Uhealth. Jan 26, 2024;12:e47843. [doi: [10.2196/47843](https://doi.org/10.2196/47843)] [Medline: [38277195](https://pubmed.ncbi.nlm.nih.gov/38277195/)]
11. Dieter V, Janssen P, Krauss I. Efficacy of the mHealth-based exercise intervention re.flex for patients with knee osteoarthritis: pilot randomized controlled trial. JMIR Mhealth Uhealth. Sep 9, 2024;12:e54356. [doi: [10.2196/54356](https://doi.org/10.2196/54356)] [Medline: [39250181](https://pubmed.ncbi.nlm.nih.gov/39250181/)]

12. Özden F, Sarı Z. The effect of mobile application-based rehabilitation in patients with total knee arthroplasty: a systematic review and meta-analysis. *Arch Gerontol Geriatr*. Oct 2023;113:105058. [doi: [10.1016/j.archger.2023.105058](https://doi.org/10.1016/j.archger.2023.105058)] [Medline: [37172329](https://pubmed.ncbi.nlm.nih.gov/37172329/)]
13. Monárez R, Mohamadi A, Drew JM, Abdeen A. Mobile application's effect on patient satisfaction and compliance in total joint arthroplasty: a systematic review and meta-analysis. *J Am Acad Orthop Surg Glob Res Rev*. Sep 1, 2023;7(9):e22.00200. [doi: [10.5435/JAAOSGlobal-D-22-00200](https://doi.org/10.5435/JAAOSGlobal-D-22-00200)] [Medline: [37678829](https://pubmed.ncbi.nlm.nih.gov/37678829/)]
14. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Serv Res*. Jan 26, 2017;17(1):88. [doi: [10.1186/s12913-017-2031-8](https://doi.org/10.1186/s12913-017-2031-8)] [Medline: [28126032](https://pubmed.ncbi.nlm.nih.gov/28126032/)]
15. Wang Q, Lee RLT, Hunter S, Zhu A, Chan SWC. Patient engagement in a mobile app-based rehabilitation program for total hip or knee arthroplasty: secondary data analysis of a randomized controlled trial. *JMIR mHealth uHealth*. Oct 1, 2024;12:e57635. [doi: [10.2196/57635](https://doi.org/10.2196/57635)] [Medline: [39353187](https://pubmed.ncbi.nlm.nih.gov/39353187/)]
16. Hong M, Loeb J, Yang M, Bailey JF. Postoperative outcomes of a digital rehabilitation program after total knee arthroplasty: retrospective, observational feasibility study. *JMIR Form Res*. Sep 19, 2022;6(9):e40703. [doi: [10.2196/40703](https://doi.org/10.2196/40703)] [Medline: [36121690](https://pubmed.ncbi.nlm.nih.gov/36121690/)]
17. Wang Q, Gao J, Zhu A, Cheng C, Chan SWC. Patient engagement in mobile technology-based rehabilitation for arthroplasty: a scoping review. *J Clin Nurs*. Sep 2025;34(9):3489-3503. [doi: [10.1111/jocn.17802](https://doi.org/10.1111/jocn.17802)] [Medline: [40346770](https://pubmed.ncbi.nlm.nih.gov/40346770/)]
18. Ramkumar PN, Haeberle HS, Ramanathan D, et al. Remote patient monitoring using mobile health for total knee arthroplasty: validation of a wearable and machine learning-based surveillance platform. *J Arthroplasty*. Oct 2019;34(10):2253-2259. [doi: [10.1016/j.arth.2019.05.021](https://doi.org/10.1016/j.arth.2019.05.021)] [Medline: [31128890](https://pubmed.ncbi.nlm.nih.gov/31128890/)]
19. Booth MW, Riegler V, King JS, Barrack RL, Hannon CP. Patients' perceptions of remote monitoring and app-based rehabilitation programs: a comparison of total hip and knee arthroplasty. *J Arthroplasty*. Jul 2023;38(7S):S39-S43. [doi: [10.1016/j.arth.2023.04.032](https://doi.org/10.1016/j.arth.2023.04.032)] [Medline: [37088222](https://pubmed.ncbi.nlm.nih.gov/37088222/)]
20. Wang Q, Lee RLT, Hunter S, Chan SWC. Patients' experiences of using a mobile application-based rehabilitation programme after total hip or knee arthroplasty: a qualitative descriptive study. *BMC Nurs*. Jul 27, 2023;22(1):246. [doi: [10.1186/s12912-023-01409-3](https://doi.org/10.1186/s12912-023-01409-3)] [Medline: [37496003](https://pubmed.ncbi.nlm.nih.gov/37496003/)]
21. Perski O, Short CE. Acceptability of digital health interventions: embracing the complexity. *Transl Behav Med*. Jul 29, 2021;11(7):1473-1480. [doi: [10.1093/tbm/ibab048](https://doi.org/10.1093/tbm/ibab048)] [Medline: [33963864](https://pubmed.ncbi.nlm.nih.gov/33963864/)]
22. MyMobility® with Apple Watch. Zimmer Biomet. 2023. URL: <https://www.zimmerbiomet.com/en/products-and-solutions/zb-edge/mymobility.html#01-overview> [Accessed 2025-12-09]
23. MyMobility® Patient App. Google Play. URL: <https://play.google.com/store/apps/details?id=com.mymobility> [Accessed 2025-12-09]
24. Privacy notice. Zimmer Biomet. 2023. URL: <https://www.zimmerbiomet.com/en/corporate/privacy-notice/privacy-notice-english.html> [Accessed 2025-12-09]
25. PIPEDA requirements in brief. Office of the Privacy Commissioner of Canada. URL: [https://www.priv.gc.ca/en/privacy-topics/privacy-laws-in-canada/the-personal-information-protection-and-electronic-documents-act-pipeda/pipeda\\_brief/](https://www.priv.gc.ca/en/privacy-topics/privacy-laws-in-canada/the-personal-information-protection-and-electronic-documents-act-pipeda/pipeda_brief/) [Accessed 2025-12-09]
26. Sekhon M, Cartwright M, Francis JJ. Development of a theory-informed questionnaire to assess the acceptability of healthcare interventions. *BMC Health Serv Res*. Mar 1, 2022;22(1):279. [doi: [10.1186/s12913-022-07577-3](https://doi.org/10.1186/s12913-022-07577-3)] [Medline: [35232455](https://pubmed.ncbi.nlm.nih.gov/35232455/)]
27. Lund AM. Measuring usability with the USE questionnaire. *Usability Interface*. 2001;8(2):3-6. URL: [https://www.researchgate.net/publication/230786746\\_Measuring\\_Usability\\_with\\_the\\_USE\\_Questionnaire](https://www.researchgate.net/publication/230786746_Measuring_Usability_with_the_USE_Questionnaire) [Accessed 2026-01-16]
28. Dawson J, Fitzpatrick R, Carr A, Murray D. Questionnaire on the perceptions of patients about total hip replacement. *J Bone Joint Surg Br*. Mar 1996;78(2):185-190. [doi: [10.1302/0301-620X.78B2.0780185](https://doi.org/10.1302/0301-620X.78B2.0780185)] [Medline: [8666621](https://pubmed.ncbi.nlm.nih.gov/8666621/)]
29. Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br*. Jan 1998;80-B(1):63-69. [doi: [10.1302/0301-620X.80B1.0800063](https://doi.org/10.1302/0301-620X.80B1.0800063)]
30. Kersten P, White PJ, Tennant A. Is the pain visual analogue scale linear and responsive to change? An exploration using Rasch analysis. *PLoS ONE*. 2014;9(6):e99485. [doi: [10.1371/journal.pone.0099485](https://doi.org/10.1371/journal.pone.0099485)] [Medline: [24921952](https://pubmed.ncbi.nlm.nih.gov/24921952/)]
31. Nuevo M, Modrego A, Rodríguez-Rodríguez D, et al. The influence of sociodemographic and health factors on adherence to home-based rehabilitation after fast-track total knee arthroplasty: secondary analysis of a randomized controlled trial. *Disabil Rehabil*. Sep 2025;47(18):4728-4737. [doi: [10.1080/09638288.2025.2458191](https://doi.org/10.1080/09638288.2025.2458191)] [Medline: [39879639](https://pubmed.ncbi.nlm.nih.gov/39879639/)]
32. Lee CMY, Thomas E, Norman R, et al. Educational attainment and willingness to use technology for health and to share health information - the reimaging healthcare survey. *Int J Med Inform*. Aug 2022;164:104803. [doi: [10.1016/j.ijmedinf.2022.104803](https://doi.org/10.1016/j.ijmedinf.2022.104803)] [Medline: [35644052](https://pubmed.ncbi.nlm.nih.gov/35644052/)]

33. Bourne RB, Chesworth B, Davis A, Mahomed N, Charron K. Comparing patient outcomes after THA and TKA: is there a difference? *Clin Orthop Relat Res*. Feb 2010;468(2):542-546. [doi: [10.1007/s11999-009-1046-9](https://doi.org/10.1007/s11999-009-1046-9)] [Medline: [19760472](https://pubmed.ncbi.nlm.nih.gov/19760472/)]

## Abbreviations

**mHealth:** mobile health  
**OHS:** Oxford Hip Score  
**OKS:** Oxford Knee Score  
**PROM:** patient-reported outcome measure  
**TFA:** theoretical framework for acceptability  
**THA:** total hip arthroplasty  
**TKA:** total knee arthroplasty  
**USE:** Usefulness, Satisfaction, and Ease of Use  
**VAS:** visual analog scale

Edited by Andre Kushniruk; peer-reviewed by Wojciech Glinkowski, Xinyuan Zhang; submitted 29.Jun.2025; final revised version received 23.Oct.2025; accepted 20.Nov.2025; published 27.Jan.2026

*Please cite as:*

Louni Y, Laroche M, Alnasser A, Abuhaneya M, Belzile E, Baskaran S, Mutch J, Albers A  
*Acceptability of a Digital Care App in Patients Undergoing Hip and Knee Arthroplasty: Prospective Cohort Study*  
*JMIR Hum Factors* 2026;13:e79682  
URL: <https://humanfactors.jmir.org/2026/1/e79682>  
doi: [10.2196/79682](https://doi.org/10.2196/79682)

© Yacine Louni, Matthew Laroche, Abdulrhman Alnasser, Mohammad Abuhaneya, Eric Belzile, Sandhya Baskaran, Jennifer Mutch, Anthony Albers. Originally published in JMIR Human Factors (<https://humanfactors.jmir.org>), 27.Jan.2026. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Human Factors, is properly cited. The complete bibliographic information, a link to the original publication on <https://humanfactors.jmir.org>, as well as this copyright and license information must be included.